

- Phan, L. T. *et al.* (2020) 'Importation and Human-to-Human Transmission of a Novel Coronavirus in Vietnam', *New England Journal of Medicine*, 382(9), pp. 872–874. doi: 10.1056/nejmc2001272.
- Ren, L. L. *et al.* (2020) 'Identification of a novel coronavirus causing severe pneumonia in human: a descriptive study', *Chinese medical journal*, 133(9), pp. 1015–1024. doi: 10.1097/CM9.0000000000000722.
- Wang, W., Tang, J. and Wei, F. (2020) 'Updated understanding of the outbreak of 2019 novel coronavirus (2019-nCoV) in Wuhan, China', *Journal of Medical Virology*, 92(4), pp. 441–447. doi: 10.1002/jmv.25689.
- Yae, Y. *et al.* (1991) 'Isolation and characterization of a thermolabile β -2 macroglycoprotein ("thermolabile substance" or 'Hakata antigen') detected by precipitating (auto) antibody in sera of patients with systemic lupus erythematosus', *Biochimica et Biophysica Acta (BBA)/Protein Structure and Molecular*, 1078(3), pp. 369–376. doi: 10.1016/0167-4838(91)90158-V.
- Zhang, X. L. and Ali, M. A. M. (2008) 'Ficolins: Structure, function and associated diseases', *Advances in Experimental Medicine and Biology*, 632, pp. 105–115. doi: 10.1007/978-0-387-78952-1_9.
- Zhu, J. *et al.* (2020) 'Clinicopathological characteristics of 8697 patients with COVID-19 in China: A meta-analysis', *Family Medicine and Community Health*, 8(2), pp. 1–11. doi: 10.1136/fmch-2020-000406.