















- poly vinyl alcohol addition and photocrosslinking on starch based materials mechanical properties,” *Carbohydr. Polym.*, vol. 60, no. 2, pp. 185–192, 2005.
- [12] I. Angelidaki *et al.*, “Defining the biomethane potential (BMP) of solid organic wastes: a proposed protocol for batch assays,” 2008.
- [13] G. K. Anderson and G. Yang, “Determination of bicarbonate and total volatile acid concentration in anaerobic digesters using a simple titration,” *Water Environ. Res.*, vol. 64, no. 1, pp. 53–59, 1992.
- [14] I. H. Franke-Whittle, A. Walter, C. Ebner, and H. Insam, “Investigation into the effect of high concentrations of volatile fatty acids in anaerobic digestion on methanogenic communities,” *Waste Manag.*, vol. 34, no. 11, pp. 2080–2089, 2014, doi: 10.1016/j.wasman.2014.07.020.
- [15] B. Budiyono, I. Syaichurrozi, and S. Sumardiono, “Biogas production from bioethanol waste: the effect of pH and urea addition to biogas production rate. Waste Technology, 1(1), 1–5. <https://doi.org/10.12777/wastech.1.1.2013.1-5>,” *Waste Technol.*, vol. 1, no. 1, pp. 1–5, 2013, doi: 10.12777/wastech.1.1.2013.1-5.