

- Shrivastava, S., and Pandey, A. K.(1981). Length-weight relationship and condition factor of three Indian major carps in composite fish farming. *Matsya*, 7: 70-74.
- Siddique, M.A.M., Khan, M.S.K., Habib, A., Bhuiyan, M.K.A., Aftabuddin, S.(2016). Size frequency and length-weight relationships of three semi-tropical cephalopods, Indian squid *Photololigo duvaucelii*, needle cuttlefish *Sepia aculeata*, and spineless cuttlefish *Sepiella inermis* from the coastal waters of Bangladesh, bay of Bengal. *Zool Ecol* 26(3):176–180
- Soni, D. D., and Kathal, K. M.(1979). Length-weight relationship in *Cirrhinus mrigala* (Val.) and *Cyprinus carpio* (Hamilton). *Matsya*, 5: 69-72.
- Stergiou, K. I., and Moutopoulos, D. K.(2001). A review of length-weight relationships of fishes from Greek marine waters. *NAGA, ICLARM Quart.*, 24(1-2): 23-39.
- Sunder, S., Kumar, K., Raina, H.S.(1984). Food and feeding habits and length-weight relationship of *Cyprinus carpio* var. *specularis* of Dal Lake, Kashmir. *Indian. J Fish.* 31(1):90-99.
- Talwar, P. K., Jhingran, A. G.(1991) *Inland fishes of India and adjacent countries* (Vols. 1&2). Oxford and IBH Publishing Co. Pvt. Ltd., New Delhi, pp. 1158.
- Ujjania, N.C., Kohli, M.P.S. and Sharma, L.L.(2012). Length-weight relationship and condition factors of Indian major carps (*Catla catla*, *Labeo rohita*, and *Cirrhinus mrigala*) in Mahi Bajaj Sagar, India. *Research Journal of Biology*, 2(1), 30-36.
- Weatherly, A.H.(1972). *Growth and ecology of the fish population*. Academic Press, London.
- www.iucnred.list.(2017) 04 February.
- Zaydın, O. O., Uckun, D., Akalın, S., Leblebici, S., and Tosunog˘lu, Z.(2007).Length-weight relationships of fishes captured from Izmir Bay, Central Aegean Sea. *J. Appl. Ichthyol.* 23 (2007), 695–696. DOI: 10.1111/j.1439-0426.2007.00853.