

10. Obasuyi BI, Fyनेface-Ogan S, Mato CN. A comparison of the haemodynamic effects of lateral and sitting positions during induction of spinal anaesthesia for caesarean section. *Int J Obstet Anesth.* 2013;**22**(2):124–8. doi: 10.1016/j.ijoa.2012.12.005. [PubMed: 23453465].
11. Patel M, Samsoun G, Swami A, Morgan B. Posture and the spread of hyperbaric bupivacaine in parturients using the combined spinal epidural technique. *Can J Anaesth* 1993; 40: 943-6.
12. Inglis A, Daniel M, McGrady E. Maternal position during induction of spinal anaesthesia for caesarean section. A comparison of right lateral and sitting positions. *Anaesthesia* 1995;50: 363-5.
13. Hajian P, Nikooseresht M, Lotfi T. Comparison of 1- and 2-Minute Sitting Positions Versus Immediately Lying Down on Hemodynamic Variables After Spinal Anesthesia with Hyperbaric Bupivacaine in Elective Cesarean Section. *Anesthesiology and Pain Medicine.* 2017;7(2):e43462. doi:10.5812/aapm.43462.
14. Kohler F, Sorensen JF, Helbo-Hansen HS. Effect of delayed supine positioning after induction of spinal anaesthesia for caesarean section. *Acta Anaesthesiol Scand.* 2002;**46**(4):441–6. [PubMed: 11952447].
15. Gori F, Corradetti F, Cerotto V, Peduto VA. Influence of positioning on plain levobupivacaine spinal anesthesia in cesarean section. *Anesthesiol Res Pract.* 2010;**2010** doi: 10.1155/2010/212696. [PubMed:20700432].
16. Obasuyi BI, Fyनेface-Ogan S, Mato CN. A comparison of the haemodynamic effects of lateral and sitting positions during induction of spinal anaesthesia for caesarean section. *Int J Obstet Anesth.* 2013 Apr;**22**(2):124-8.