





















49. Iqbal S, Zia-Ul-Haq M, Ahmad S; (2007). Nutritional and Compositional study of Desi chickpea (*Cicer arietinum* L.) cultivars grown in Punjab, Pakistan; Food Chemistry. 105; 1357-1363.
50. GS Ranhotra, BK Glaser et.al., (1993). Composition and protein nutritional quality of quinoa, cereal chemistry, 70, 303-303.
51. KJ Lorenz, GS Ranhotra, JA Gelroth, BK Glaser, DL Johnson; Composition in protein, nutritional quality of Quinoa; Cereal Chemistry; 1993; 70; 303.
52. Atiya Techaparin, Kimroeun, Vann. (2020). Beans germination as potential tool for GABA – enriched tofu production, Food science and technology, 57(11), 3947- 3954.
53. Sandeep Kumar, Mukul Das, (2013), Clinical complications of kidney bean consumption, Nutrition 29(6), 821- 827.
54. PV Wagh, IE Liener. (1963). Nutritive value of red kidney bean for chicks, The journal of nutrition 80 (2), 191-195.
55. Dina anwar, el- chaghaby Ghadir , (2019). Nutritional quality, amino acid profiles , protein digestibility corrected amino acid and anti-oxidant properties of fried tofu and seitan , Food and environment safety journal, 18 (3).
56. Vijay Jayasena, WS Khu, et.al, (2010). The development and sensory acceptability of lupin based tofu, The journal of food quality, 33 (1), 85-97.