

- [5] Sorenson RL, Brelje TC. Prolactin receptors are critical to the adaptation of islets to pregnancy. *Endocrinology* (2009); 150(4): 1566–9.
- [6] Huang C, Snider F, Cross JC. Prolactin receptor is required for normal glucose homeostasis and modulation of beta-cell mass during pregnancy. *Endocrinology* (2009); 150(4): 1618–26.
- [7] Clare Tower: Gestational diabetes In: David M.L and mark D.k evidence – based text for the MRCOG 3rd edition (2016); 25: 192.
- [8] Sara N. Ali and Anne Dornhorst. Diabetes in Pregnancy. Dewhurst' text book of Obstetrics and Gynecology. 9th edition (2018); (9) 99 -100.
- [9] Gargee Pandit and Kaizad R. diabetes in pregnancy Practical Guide to High. Risk Pregnancy and Delivery. Fourth Edition (2015) chapter 15; page 257.
- [10] EI Ekinci, N Torkamani, SK Ramchand, et al. Higher maternal serum prolactin levels are associated with reduced glucose tolerance during pregnancy, *Journal of diabetes* (2017); p 697-700.
- [11] Shalayel MH, Elrobb MS, Idris SA, Mohammed MS, Ahmed SA. Prolactin and insulin estimates in pregnancy with glucose intolerance. *Pakistan Journal of Medical Science* (2010); 26(1): 102-6.
- [12] Daimon M, Kamba A, Murakami H, et al. Association between serum prolactin levels and insulin resistance in non-diabetic men (2017).
- [13] Tiange Wang, Jieli Lu, Yu Xu et al. Circulating Prolactin Associates With Diabetes and Impaired Glucose Regulation, *Diabetes Care* (2013); 36(7): 1974–1980.
- [14] Deepti Shroff Karhade and David C. Wang, Intrapartum serum prolactin may predict risk of postpartum diabetes, June 25(2016).
- [15] Ravi Retnakaran, Chang Ye, Caroline K. Kramer, et al. Maternal Serum Prolactin and Prediction of Postpartum β -Cell Function and Risk of Prediabetes/Diabetes (2016).
- [16] M.A. Ramos-Roman, Prolactin and Lactation as Modifiers of Diabetes Risk in Gestational Diabetes (2011); 43(09): 593-600.