Effects of Probiotics and Herbal oils in Reducing Multidrug Resistant *E. coli* Infection and Promoting Growth in Rabbits

Mona S. Abdallah¹*, Heba M. Hassan², Wafaa A.A Ibrahim³, Amira M. Helal³, and Engy A. Hamed²

¹Department of Avian and Rabbit Medicine, Faculty of Veterinary Medicine, Suez Canal University, Ismailia, Egypt 41522. ²Reference Laboratory for Veterinary Quality Control on Poultry production, Animal Health

Research Institute, Agricultural Research Center (ARC). Nadi El-Seid Street, Dokki P.O. Box 246, Giza 12618, Egypt.

³Biotechnology Department, Reference Laboratory for Veterinary Quality Control on Poultry production, Animal Health Research Institute, Agricultural Research Center (ARC). Ismailia, Egypt

* Corresponding author: Mona S. Abdallah, Department of Avian and Rabbit Medicine, Faculty

of Veterinary Medicine, Suez Canal University, Ismailia, Egypt. E-mail address: <u>mona_salem@vet.suez.edu.eg</u> Address: Kilo 4.5 Ring Road, Ismailia, 41522, Egypt.

ABSTRACT

Colibacillosis is a common infectious bacterial disease that can cause enteritis and high mortality in young rabbits, threatening the breeding industry. This work performed to express some virulence-determining genes and antimicrobial resistance patterns of isolated E. coli and to study the effect of probiotics and/or herbal oils on E. coli infection in rabbits. Collection of vaginal swabs and semen samples from rabbits with reproductive problems (farm 1) and anal swabs from rabbits with diarrhea (farm 2) in Ismailia Governorate, Egypt. Samples were subjected for bacterial identification, antibiogram-testing and molecular monitoring of iss and eaeA virulence genes. An experiment was done, 54 weaned California rabbits were divided into 6 equal groups, negative control (G1), non-infected treated with probiotic and both fennel and moringa oils (G2), Infected and treated with probiotic, fennel and moringa oils (G3), treated with probiotic only (G4), treated with colistin sulfate (G5) and Positive control (G6). Infected groups (G3, G4, G5 and G6) were challenged orally with E. coli ($1x10^7$ CFU). The detection of E. coli was 15.3% (2/13) and 62.5% (10/16) in farm (1) and (2) respectively. The isolated E. coli was multidrug-resistant (MDR), carrying virulence genes iss and eaeA, with prevalence of 100% (12/12) and 50% (6/12) respectively. Clinical symptoms were prominent in the experiment E. coli infected untreated group (6), with the least evidence in group (5). The E. coli colony counts were significantly higher in group (6) while rabbits in group (3) had significantly higher body weights in comparison to the other infected groups. Thus we concluded that, the use of two herbal oils and probiotics were able to reduce mortality rates, clinical signs, and the total E. coli count in experimentally infected rabbits.

KEY WORDS: E. coli, herbal oils, Probiotics, Rabbits, virulence gene

INTRODUCTION

Rabbits are an important source of animal protein for many countries in the world (El-Deep et al., 2021). Rabbits' meat has a healthy source of protein with low fat and cholesterol especially for young and old people (Imam et al., 2020). Therefore, in order to maintain the growth and development of rabbits, optimal growth conditions should be maintained. (El-Deep et al., 2020a).

Young rabbits are particularly susceptible to various diseases such as gastrointestinal disorder especial during the weaning period, and the main pathogenic bacteria is *Escherichia coli* (*E. coli*) which cause high economic loss in rabbits farming, It can be transmitted to rabbits through ingestion of contaminated food and water, causing loss of appetite, weakness, diarrhea, slowness of movement, crouching posture, abdominal distention and death (Camer et al., 2012).

Virulence-determining genes influence the pathogenicity of bacteria in the host and allow them to produce enteric or extraintestinal disease, the *eae* gene is considered the main genetic marker used to describe enteropathogenic *E. coli* strains (Milon et al., 1999), enable bacteria to adhere to the epithelial cells in intestinal lumen (Badagliacca et al., 2018). Episomal increased serum survival protein (*iss*) is a good indicator of the pathogenicity of bacteria and is more common in infective strains (Pfaff-McDonough et al., 2000).

Rabbit breeders often use antibiotics to treat and prevent such diseases, but the continuous use of these antibiotics may harm the natural immunity of animals, and accumulate in meat, which could transmit to the human and weak the natural immunity against infection (El-Deep et al., 2020b). Prohibition of antibiotics and discovery of new alternative biological additives forced scientists to study some non-antibiotic compounds such as prebiotics, probiotics, herbal extracts, and organic acids (Oso et al., 2013; Olorunsola et al., 2016).

Probiotics play many roles in rabbits farming, like decreasing the incidence of gastrointestinal infection and increase immune response which may help in decreasing use of antibiotics, enhancing of rabbits growth due to increase the feed conversion rates which lead to increase the quality of final product (Mancini and Paci, 2021). Combination of medicinal herbs and their extracts primarily used as natural immune boosters, growth promotion and antioxidant supply to promote animal health and productivity (Al-Sagan et al., 2020; Dawood, 2021).

As one of the most promising alternative feedstuffs, *Moringa oleifera* (moringa) has antioxidant properties and strong antimicrobial activity, and can be used as a natural food antibacterial agent in food systems (Sharma et al., 2020). As well as *fennel Foeniculum* (fennel), which has antimicrobial properties and may be used to reduce the number of undesirable gut microbes and enhance digestion (Elgayyar et al., 2001).

The current study aimed to monitor the antimicrobial resistance patterns and the presence of certain virulence-determining genes (*iss* and *eaeA*) in *Escherichia coli* isolated from rabbits. As well as investigating the efficacy of fennel, moringa oil, and/or probiotics as alternatives to antibiotics in reducing *E. coli* infections and promoting growth in rabbits.

MATERIAL AND METHODS

Ethical approval

The materials and protocols applied in this scientific study were accepted by the Scientific Research Ethics Committee of the Faculty of Veterinary Medicine, University of Suez, Ismailia, Egypt.

Sampling and clinical examination

Twenty-nine samples; Vaginal swabs (n=9), semen samples (n=4), anal swabs (n=12) and liver specimens (n=4) were collected under sterile conditions from two rabbit farms in Ismailia province, Egypt. Vaginal swabs were gathered from rabbit's suffering from reproductive problems (infertility, abortion, dystocia, and stillbirth) while semen samples were collected from animals with history of sperm abnormalities farm (1). Anal swabs and liver specimens were collected from clinically diarrhetic rabbits (farm 2) with dirty fur around anal opening, and combined with weight loss, lethargy, and anorexia. The obtained samples were immediately inoculated on peptone water.

E. coli isolation and identification

Loops of peptone water were inoculated aseptically in nutrient broth (Oxoid, UK) and incubate aerobically at 37°C for 24-48 h. Eosin methylene blue (EMB) and MacConkey agar (Oxoid, UK) were streaked using loopful of broth-culture. Identification of isolated colonies was done based on their colonial characters, microscopical examination, and biochemical reactions (indole, oxidase, Voges-Proskauer, methyl-red, lactose fermentation, H2S, citrate-utilization, and urease tests) as mentioned by (Quinn et al., 2011).

Antimicrobial susceptibility testing

Escherichia coli isolates were monitored for susceptibility to antimicrobial agents using the disk diffusion method. Using these antimicrobials; ampicillin (AMP 10 μ g), tetracycline (T 30 μ g), norfloxacin (NOR 10 μ g), streptomycin (S 10 μ g), fosfomycin (Fo 200 μ g), colistin (CL 10 μ g), penicillin (P 10u), erythromycin (E 15 μ g) and lincospectin (LS 100 μ g) (Oxoid, UK). The sensitivity degree was interpreted after incubating the plates at 37°C for 18 h according to (CLSI, 2017).

Molecular typing of E. coli virulence-determining genes

Monitoring of *E. coli* virulence-determinant genes (*iss* and *eaeA*) was performed. These genes were selected based on their important roles in disease induction as mentioned in previous studies (Hassan and Al-Azeem, 2009; Eldin and Reda, 2016). DNA extracted from the examined samples using the QIAamp DNA Mini kit (Qiagen, Germany, GmbH) with modifications as for the manufacturer's instruction. Reaction volume was 25-µl and contained 1 µl of each primer with concentration of 20 pmol, 12.5 µl of Emerald Amp Max PCR Master Mix (Takara, Japan), 5 µl of DNA template, and 5.5 µl of water. Performing the reaction was in an Applied biosystems 2720 thermal cycler. Primers used were obtained from Metabion (Germany) and are listed in table (1). The separation of PCR products was applied by electrophoresis on 1.5% agarose gel stained with 0.5µg/ml ethidium bromide (Applichem, Germany, GmbH). Gel documentation system (Alpha Innotech, Biometra) was used to photograph the gels, and the data were evaluated by computer software.

Experimental design

This experiment was designed to evaluate *in vitro* and/or *in vivo* antimicrobial action of herbal oils and probiotic against *E. coli* field isolate. The experiment was done in experimental Research Center, Animal Health Research Institute, Dokki, Giza, Egypt

Bacterial culture

E. coli strain used in this experiment was obtained from field isolates in this study after bacteriologically, serologically, and molecularly identification of virulence gene. The *E. coli* strain was inoculated in peptone water and incubated for 24 h at 37°C. Then prepare an inoculum of 10^7 and 10^9 colony forming units (CFU)/ml based on Mc-Farland criteria.

Drug used in this experiment

Colistin Sulfate: (Royal link Pharma) On 3rd day post-infection (PI) a dose of 20 mg/kg bwt was administered orally for 5 days.

ProBax probiotic: (Key Vet Cooperation KVC) contains (*Lactobacillus Casei* $1x10^{11}$ CFU and *Bacillus subtillus* $1x10^{11}$ CFU). Dosing at 1 gm/L in drinking water throughout the experiment. Fennel and moringa oils were optioned from National Research Center, Egypt. Giving in drinking water at dose of 6ml/L throughout the experiment period.

Antimicrobial activity of herbal oil (in vitro test)

The *in vitro* antibacterial effects of fennel and moringa oils were assessed and appropriate levels of these oils against challenged *E. coli* strain was determined. Antibacterial evaluation was performed according to the MIC assay method according (Kowalska-Krochmal and Dudek-Wicher, 2021). Mueller Hinton Broth (100 μ l) was Poured into wells of a microtiter plate, then 100 μ l of herbal oil was added to the first well, followed by a double-fold serial dilution. Next, 100 μ l of E. coli field isolate suspension (109 cfu/ml), estimated at 0.5 Mc-Farland, was inoculated into each well and incubated at 37 °C for 24 h. Bacterial growth was indicating by existence of turbidity. MIC is the smallest concentration of herbal oil abled to inhibit the bacterial growth after incubation for 24- 48 h.

In vivo evaluation of the effectiveness of probiotics and herbal oils in rabbits experimentally infected with E. coli

Fifty-eight weaned California rabbits (30 days old) of both sexes were purchased from a private rabbit's farm in Cairo, Egypt. On arrival, four rabbits were sacrificed, and their livers cultured to confirm that they were free of *E. coli*. During the experiment, the rabbits were kept in metal cages with good sanitary conditions and fed commercial balanced diets ration ad libitum. Rabbits were randomly divided into 6 equal groups (9 rabbits each), G1: Negative control: non infected non treated. G2: Non infected and treated with probiotic (at dose rate of 1 ml/L in drinking water) and both fennel and moringa oils (at dosage rate of 6 ml/L in drinking water) throughout the experiment period. G3: Infected and treated with probiotic (at dose rate of 1 ml/L in drinking water) and both fennel and moringa oils (at dosage rate of 6 ml/L in drinking water) throughout the experiment period. G4: Infected and treated with probiotic only (at dose rate of 1 ml/L in drinking water). G5: Infected and treated with colistin sulfate orally at the 3rd day post-infection (at dosage rate of 20 mg/kg bwt. for 5 consecutive days). G6: Positive control: infected

non treated. Infected groups were orally administered 0.1 ml of *E. coli*, the infectious dose $(1 \times 10^7 \text{ CFU})$.

Mortality rate, clinical signs, post-mortem examination, and weekly mean body weight were recorded during the experiment period (1 month)

Total bacterial counts

Total *E. coli* counts of the infected groups at 7th, 15th, and 21st day PI was evaluated. One gram of liver tissue was added into 9 ml sterile nutrient broth and homogenized well later the suspension was vortexed. Serial 10- fold dilutions were made. One ml of each dilution was inoculated onto double MacConkey agar plates and incubated aerobically at $37C^{0}$ for 24-48 h and the number of CFU per gram of liver was determined.

Statistical analysis

Statistical analysis and graphs were performed with R (A language and environment for statistical computing) (R Core Team, 2020). Results were described as mean \pm SD, comparison of treatments for each group was evaluated by analysis of variance (ANOVA), then for multiple comparisons the Tukey's post-hoc test was applied, and calculation of p-value.

RESULTS

Detection of E. coli in tested samples

The identification of isolated *E. coli* was done, based on their morphology and biochemical reaction. *E. coli* grown on MacConkey and EMB agar to produce characteristically growth colonies, pink and metallic sheen colonies, respectively, which appear microscopically as medium-sized Gram-negative bacilli. Biochemically, *E. coli* isolates tested positive for indole, methyl red, and lactose fermentation. Additionally, they tested negative for urease, Voges-Proskauer, H2S production and citrate utilization. Detection of *E. coli* was 22.2% (2/9), 0% (0/4), 66.6% (8/12), and 50% (2/4) from vaginal swabs, semen samples, anal swabs, and liver samples respectively, for an overall occurrence of 15.3% (2/13) and 62.5% (10/16) in farm (1) and (2) respectively.

Antimicrobial resistance pattern of isolated E. coli

Antimicrobial susceptibility tests revealed that *E. coli* isolates from both farms were only sensitive to three antibiotics, norfloxacin, fosfomycin and colistin, while *E. coli* isolated from farm (2) was intermediate sensitive to ampicillin and streptomycin. As well as the examined isolates displayed high resistance to erythromycin, lincospectin, penicillin, and oxytetracycline besides, streptomycin in farm (1) as shown in Table 2.

Detection of virulence-determining genes of E. coli isolates

Molecular detection of *E. coli* virulence-determining genes showed that the corrected size of the *iss* positive DNA band was 266 bp (Figure 1a), and the corrected size of the *eaeA* gene

was 248 bp (Figure 1b). The *iss* and *eaeA* virulence genes were present in 100% (12/12) and 50% (6/12) of the tested isolates, respectively as shown in Figure (1).

In vitro assessment of the antimicrobial activity of herbal oil against E. coli isolate

Antimicrobial activity of fennel and moringa oils against isolated *E. coli* was evaluated by determination of MIC. From the results obtained, it was noted that *E. coli* was sensitive to both oils. These oils had MIC value of 1:16 and were able to completely inhibit the growth of bacteria on MacConkey agar after incubation for 24 h at 37° C.

In vivo evaluation of the effectiveness of probiotics and herbal oils against E. coli infection in rabbits

Clinical symptoms, mortality rate and post-mortem examination

Symptoms appeared on the 3rd day PI, infected rabbits suffered from anorexia, general depression, dullness, abdominal distention, perineal staining with watery, brown diarrhea. These findings were prominent in the control-infected untreated group (6), followed by the probiotic treated- infected group (4), with the least evidence in the colistin sulfate-treated group (5). Control positive (G6) record the highest mortality rate during the experiment, which was 33.3% (3/9), while that of the other infected groups was 11.1% (1/9). Dead rabbits showed congestion in all internal organs, pericarditis, enteritis, ballooning of the intestine with petechial hemorrhage on intestinal wall.

Effects of probiotics, herbal oils and colistin on the total colony count of E. Coli in experimentally infected rabbits

E. coli experimentally infected and untreated rabbits (G6) showed significantly higher total colony counts compared to the other infected-treated groups (3, 4, and 5) during experiment period. Whereas groups (3, 4 and 5) recorded nonexistence of *E. coli* counts at the 21^{st} PI, as shown in Figure (2).

Effects of probiotics, herbal oils and colistin on body weight of rabbits

The effect of commercial probiotic and two herbal oils and colistin antibiotic on body weight of healthy and *E. coli* experimentally infected rabbits at 7th ,15th and 21st days PI was illustrated in Table (3). At 7th day experimentally infected rabbits with *E. coli* and probiotic treated (G4) and colistin treated (G5) showed body weights significantly decreased as compared to other groups. At 15th and 21st day PI experimentally infected rabbits with *E. coli* and treated with probiotic and both fennel and moringa oils (G3) showed body weights significantly increased as compared to other groups. In non-infected groups rabbits treated with both fennel and moringa oils and probiotic (G2) showed body weights significantly increased at 7th ,15th and 21st day of experiment in comparison to non-treated group (1).

DISCUSSION

Colibacillosis, is a common and major infectious disease of rabbits that endangers the rabbit industry (Zhao et al., 2018). The virulence factors associated with the pathogenicity of E.

coli infection are several and display a broad range of activities, including toxins, iron acquisition, and adhesion factors (Sarowska et al., 2019). The aim of this study was to monitor the presence of two important virulence-determining genes (*iss* and *eaeA*) and the pattern of antimicrobial resistance in *Escherichia* coli isolated from rabbits as well as the efficacy of herbal oil and or probiotic as an antibiotic alternative compared to colistin sulfate for reducing *E. coli* infection and promoting growth in Rabbits.

The Bacteriological examination revealed that the detection of *E. coli* was 15.3% (2/13) and 62.5% (10/16) in farm 1 (suffering from reproductive problems) and farm 2 (clinically diarrhetic rabbits) respectively. *Escherichia coli* is a commensal or pathogenic bacterium that can cause intestinal or extraintestinal infection in rabbits (Okerman, 1994). There was a variation in the detection of *E. coli* among the inspected two farms, but it is difficult to generalize this result due to the small number of samples. Our results are consistent with a previous report that *E. coli* was isolated from 70.12% of diarrheal rabbits (Sakr et al., 2019) as well as 20% and 16.66% of non-aborted does and aborted live does respectively (EL-Sayed and Abd EL-latif, 2006).

Combination of virulence-determining genes and antibiotic resistance genes may lead to emergence of unexpected new bacterial strains with high mortality and morbidity. In this study, the *iss* and *eaeA* virulence genes were present in 100% (12/12) and 50% (6/12) of *E. coli* isolates detected using PCR, a finding consistent with other studies (Eid et al., 2017; Roshdy et al., 2021). With regard to antimicrobial susceptibility assessment, recovered *E. coli* isolates exhibited significant resistance to tetracycline, erythromycin, lincospectin and penicillin, which indicate *E. coli* isolated in this study are multidrug resistant (MDR) isolates (Moawad et al., 2017). Widespread use of antibiotics in an uncontrolled manner in the veterinary sector leads to the development of such resistant strains (Algammal et al., 2020b). This result is in line with (Aboelhadid et al., 2022) who mentioned that *E. coli* isolates displayed high resistance to almost all antimicrobials tested.

The efficacy of drugs in the treatment of bacterial diseases requires not only the active search for new therapeutic strategies, but also the careful selection of antibiotics based on various parameters, including antimicrobial susceptibility testing (Kowalska-Krochmal and Dudek-Wicher, 2021), Isolated *E. coli* showed susceptibility to norfloxacin, fosfomycin, and colistin sulfate. Based on antibiotic susceptibility testing and previous reports (Algammal et al., 2020a), which inform that colistin sulfate has good activity against MDR *E. coli*. A trial was performed to investigate the effectiveness of herbal oils and/or probiotics compared with colistin sulfate in reducing *E. coli* infections.

In this experiment, the rabbits infected with *E. coli* developed symptoms such as anorexia, dullness, abdominal distension, and brown watery diarrhea on the 3rd day PI. Necropsy of dead rabbit revealed congestion in all visceral organs, pericarditis, enteritis, ballooning of the intestine with petechial hemorrhage on the wall, these results are similar to previous reports (Prescott, 1978; Licois, 2004; Ismail et al., 2017). The symptoms were prominent in infected untreated group (6), followed by probiotic treated-infected group (4), with the least evidence in the colistin sulfate-treated group (5). The choice of treatment should be based on susceptibility results, colistin may be a good option against multidrug resistant (Gram-negative) bacteria (Li et al., 2005).

In terms of mortality, the infected untreated group (6) had the highest mortality rate during the experiment, 33.3% (3/9), while the mortality rate of other infected groups was 11.1% (1/9). The use of probiotics, fennel and moringa oils, and colistin sulfate was able to reduce

mortality in this experiment, as probiotic supplementation overcame the severity of clinical symptoms and reduced mortality in experimentally infected rabbits (Abdelhady and El-Abasy, 2015). More recently moringa is commonly used in several medicinal applications to control various digestive disorders (Gupta et al., 2018) as well as there is evidence that the *fennel Foeniculum* constituents had antimicrobial properties (Mohammed and Abbas, 2009).

The total colony count of E. coli was significantly higher in the experimentally infected and untreated rabbits (G6) compared to the other groups (3, 4, and 5) during the experiment. However, groups (3, 4, and 5) had no *E. coli* counts recorded at 21st PI, this can be attributed to the dominance of gut beneficial bacteria over pathogenic bacteria or inhibitory effect of probiotics on E. coli in the gut (Mattar et al., 2001; Panda et al., 2010). In addition, rabbits in the probiotic and herbal oil treatment group (3) gained significantly more body weight compared to the other infection groups. Infections may directly reduce performance and feed efficiency by disrupting gastrointestinal absorption capacity or indirectly by reducing normal gut microflora and metabolic processes (Abou-Kassem et al., 2021). But supplementation with fennel oil improves feed conversion and body weight by increasing length of intestinal villi. (Imbabi et al., 2021). Fennel is useful for degassing and reducing gas from dietary fermentation in the gastrointestinal tract, and also contains anethole, which eliminates pathogenic bacteria and improves body weight and feed conversion (Mohammed and Abbas, 2009). Synergy of herbal oils and probiotics enhances body weight due to the dynamic effect of probiotics on competing intestinal pathogenic microorganisms and/or enhancing immunity, fighting infectious agents, promoting digestion, absorption, promoting intestinal development (Chen et al., 2018)

CONCLUSION

Escherichia coli isolates were multidrug resistant (MDR) to erythromycin, lincospectin, penicillin, and oxytetracycline, and carried the virulence genes *iss* and *eaeA* with a high prevalence. Herbal oils and probiotics synergistically enhance body weight, compete with *Escherichia coli* infection, reduce mortality and clinical symptoms in experimentally infected rabbits.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

REFERENCES

- Abdelhady, D.H., El-Abasy M.A., 2015. Effect of Prebiotic and Probiotic on Growth, Immunohematological responses and Biochemical Parameters of infected rabbits with Pasteurella multocida. Benha Veterinary Medical Journal. 28, 40-51.
- Aboelhadid, S.M., Hashem S.A., Abdel-Kafy E.-S.M., Abdel-Baki A.-A.S., Al-Quraishy S., Abed A.H., Kamel A.A., 2022. Prevalence and antimicrobial sensitivity of Escherichia coli and Salmonella species in field cases of rabbit intestinal coccidiosis treated with prebiotic. Austral journal of veterinary sciences. 54, 9-16.
- Abou-Kassem, D.E., Mahrose K.M., El-Samahy R.A., Shafi M.E., El-Saadony M.T., Abd El-Hack M.E., Emam M., El-Sharnouby M., Taha A.E., Ashour E.A., 2021. Influences of

dietary herbal blend and feed restriction on growth, carcass characteristics and gut microbiota of growing rabbits. Italian Journal of Animal Science. 20, 896-910.

- Al-Sagan, A.A., Khalil S., Hussein E.O., Attia Y.A., 2020. Effects of fennel seed powder supplementation on growth performance, carcass characteristics, meat quality, and economic efficiency of broilers under thermoneutral and chronic heat stress conditions. Animals. 10, 206.
- Algammal, A.M., Hetta H.F., Batiha G.E., Hozzein W.N., El Kazzaz W.M., Hashem H.R., Tawfik A.M., El-Tarabili R.M., 2020a. Virulence-determinants and antibiotic-resistance genes of MDR-E. coli isolated from secondary infections following FMD-outbreak in cattle. Scientific reports. 10, 1-13.
- Algammal, A.M., Mabrok M., Sivaramasamy E., Youssef F.M., Atwa M.H., El-Kholy A.W., Hetta H.F., Hozzein W.N., 2020b. Emerging MDR-Pseudomonas aeruginosa in fish commonly harbor oprL and toxA virulence genes and blaTEM, blaCTX-M, and tetA antibiotic-resistance genes. Scientific reports. 10, 1-12.
- Badagliacca, P., Agnoletti F., Drigo I., Merildi V., Lopes F., Pompilii C., Mangone I., Scacchia M., Tonelli A., Masson L., 2018. Virulence gene profiles of rabbit enteropathogenic Escherichia coli strains isolated in Northern Italy. Veterinaria Italiana. 54, 189-196.
- Bisi-Johnson, M.A., Obi C.L., Vasaikar S.D., Baba K.A., Hattori T., 2011. Molecular basis of virulence in clinical isolates of Escherichia coli and Salmonella species from a tertiary hospital in the Eastern Cape, South Africa. Gut pathogens. 3, 1-8.
- Camer, G.A., Roh Y.-S., Cho A.-R., Kim J.-W., Umanets A., Kim B.-S., Lim S.-K., Lee H.-S., Lim C.-W., 2012. A case of nonserotypable Escherichia coli infection in a Korean rabbit farm. Korean Journal of Veterinary Service. 35, 69-71.
- Chen, H.-J., Yang W.-Y., Wang C.-Y. (2018). The review on the function of intestinal flora and the regulatory effects of probiotics on the intestinal health of rabbits. Paper presented at the 2017 2nd International Conference on Biological Sciences and Technology (BST 2017).
- CLSI, C.a.L.S.I., 2017. Clinical and Laboratory Standards Institute: Performance standards for antimicrobial susceptibility testing. 27th ed 424 informational supplement. CLSI Doc. M100-S20).
- Dawood, M.A., 2021. Nutritional immunity of fish intestines: Important insights for sustainable aquaculture. Reviews in Aquaculture. 13, 642-663.
- Eid, H., Abd El Gawad N., Zaghlol E., 2017. Bacteriological studies on Escherichia coli infection in rabbits. Suez Canal Veterinary Medical Journal. SCVMJ. 22, 133-144.
- El-Deep, M.H., Amber K.A., Eid Y.Z., Alrashood S.T., Khan H.A., Sakr M.S., Dawood M.A., 2020a. The influence of dietary chicken egg lysozyme on the growth performance, blood

health, and resistance against Escherichia coli in the growing rabbits' cecum. Frontiers in Veterinary Science. 7, 579576.

- El-Deep, M.H., Amber K.A., Elgendy S., Dawood M.A., Elwakeel E.M., Paray B.A., 2020b. Oxidative stress, hemato-immunological, and intestinal morphometry changes induced by ochratoxin A in APRI rabbits and the protective role of probiotics. Environmental Science and Pollution Research. 27, 35439-35448.
- EL-Sayed, H.M., Abd EL-latif M., 2006. Studies on some Bacteria associated with abortion in rabbits. Assiut Veterinary Medical Journal. 52, 285-293.
- El-Deep, M.H., Dawood M.A., Assar M.H., Ahamad Paray B., 2021. Aspergillus awamori positively impacts the growth performance, nutrient digestibility, antioxidative activity and immune responses of growing rabbits. Veterinary Medicine and Science. 7, 226-235.
- Eldin, W.F.S., Reda L.M., 2016. Prevalence of diarrheagenic Escherichia coli in suckling rabbits. Japanese Journal of Veterinary Research. 64, S149-S153.
- Elgayyar, M., Draughon F., Golden D., Mount J., 2001. Antimicrobial activity of essential oils from plants against selected pathogenic and saprophytic microorganisms. Journal of food protection. 64, 1019-1024.
- Gupta, S., Jain R., Kachhwaha S., Kothari S., 2018. Nutritional and medicinal applications of Moringa oleifera Lam.—Review of current status and future possibilities. Journal of Herbal Medicine. 11, 1-11.
- Hassan, S., Al-Azeem M., 2009. Determination of virulence gene markers and antimicrobial resistance in Escherichia coli isolated from rabbit in Egypt. Global Veterinaria. 3, 260-267.
- Imam, M.A.A., Dorina M., Mohamed S., Ayman A., Monica M., 2020. Rabbits Meat Production in Egypt and its Impact on Food Security, Small Holders Income and Economy. Agricultural Research & Technology: Open Access Journal. 24, 81-85.
- Imbabi, T., Sabeq I., Osman A., Mahmoud K., Amer S.A., Hassan A.M., Kostomakhin N., Habashy W., Easa A.A., 2021. Impact of Fennel Essential Oil as an Antibiotic Alternative in Rabbit Diet on Antioxidant Enzymes Levels, Growth Performance, and Meat Quality. Antioxidants. 10, 1797.
- Ismail, A., Abdien H., Hamed D., El Feil W.M., 2017. Prevalence of some Enteric Bacterial Infections Causing Rabbit Enteritis and Attempts to Control Rabbit Coli Enteritis with Phytobiotics. Zagazig Veterinary Journal. 45, 91-101.
- Kowalska-Krochmal, B., Dudek-Wicher R., 2021. The minimum inhibitory concentration of antibiotics: Methods, interpretation, clinical relevance. Pathogens. 10, 165.

- Li, J., Nation R.L., Milne R.W., Turnidge J.D., Coulthard K., 2005. Evaluation of colistin as an agent against multi-resistant Gram-negative bacteria. International journal of antimicrobial agents. 25, 11-25.
- Licois, D. (2004). Domestic rabbit enteropathies. Paper presented at the Proceedings of the eighth world rabbit congress.
- Mancini, S., Paci G., 2021. Probiotics in rabbit farming: Growth performance, health status, and meat quality. Animals. 11, 3388.
- Mattar, A., Drongowski R., Coran A., Harmon C., 2001. Effect of probiotics on enterocyte bacterial translocation in vitro. Pediatric surgery international. 17, 265-268.
- Milon, A., Oswald E., De Rycke J., 1999. Rabbit EPEC: a model for the study of enteropathogenic Escherichia coli. Veterinary research. 30, 203-219.
- Moawad, A.A., Hotzel H., Awad O., Tomaso H., Neubauer H., Hafez H.M., El-Adawy H., 2017. Occurrence of Salmonella enterica and Escherichia coli in raw chicken and beef meat in northern Egypt and dissemination of their antibiotic resistance markers. Gut pathogens. 9, 1-13.
- Mohammed, A.A., Abbas R.J., 2009. The effect of using fennel seeds (Foeniculum vulgare L.) on productive performance of broiler chickens. International Journal of Poultry Science. 8, 642-644.
- Okerman, L., 1994. Diseases of the digestive system. In: Price C. J (Ed), Diseases of Domestic Rabbits. Blackwell Scientific Publications, Oxford. 71-91.
- Olorunsola, R., Akinduti P., Oso A., Akapo A., Eruvbetine D., Oyekunle M., 2016. Effect of dietary supplementation with probiotics and prebiotics on haematological indices, serum chemistry and gut Salmonella count of broilers sourced from Salmonella-infected hatcheries in South-west zone of Nigeria. Am. J. Exp. Agric. 10, 1-13.
- Oso, A., Idowu O., Haastrup A., Ajibade A., Olowonefa K., Aluko A., Ogunade I., Osho S., Bamgbose A., 2013. Growth performance, apparent nutrient digestibility, caecal fermentation, ileal morphology and caecal microflora of growing rabbits fed diets containing probiotics and prebiotics. Livestock Science. 157, 184-190.
- Panda, A., Tatarov I., Melton-Celsa A.R., Kolappaswamy K., Kriel E.H., Petkov D., Coksaygan T., Livio S., McLeod C.G., Nataro J.P., 2010. Escherichia coli O157: H7 infection in Dutch belted and New Zealand white rabbits. Comparative medicine. 60, 31-37.
- Pfaff-McDonough, S.J., Horne S.M., Giddings C.W., Ebert J.O., Doetkott C., Smith M.H., Nolan L.K., 2000. Complement resistance-related traits among Escherichia coli isolates from apparently healthy birds and birds with colibacillosis. Avian Diseases. 23-33.
- Prescott, J., 1978. Escherichia coli and diarrhoea in the rabbit. Veterinary Pathology. 15, 237-248.

- Quinn, P.J., Markey B.K., Leonard F.C., Hartigan P., Fanning S., Fitzpatrick E. (2011). Veterinary microbiology and microbial disease: John Wiley & Sons.
- Roshdy, H., Shalaby A.G., Mohamed A.A.E., Badr H., 2021. Detection of aerobic bacterial pathogens associated with early embryonic death in pregnant New Zealand female Rabbits in Egypt. Veterinary World. 14, 986.
- Sakr, M.M., Ebied S.K., Ibrahim M.S., 2019. Detection of Diarrheagenic Escherichia Coli in Rabbits. Alexandria Journal for Veterinary Sciences. 62,
- Sarowska, J., Futoma-Koloch B., Jama-Kmiecik A., Frej-Madrzak M., Ksiazczyk M., Bugla-Ploskonska G., Choroszy-Krol I., 2019. Virulence factors, prevalence and potential transmission of extraintestinal pathogenic Escherichia coli isolated from different sources: recent reports. Gut pathogens. 11, 1-16.
- Sharma, P., Wichaphon J., Klangpetch W., 2020. Antimicrobial and antioxidant activities of defatted Moringa oleifera seed meal extract obtained by ultrasound-assisted extraction and application as a natural antimicrobial coating for raw chicken sausages. International Journal of Food Microbiology. 332, 108770.
- Yaguchi, K., Ogitani T., Osawa R., Kawano M., Kokumai N., Kaneshige T., Noro T., Masubuchi K., Shimizu Y., 2007. Virulence factors of avian pathogenic Escherichia coli strains isolated from chickens with colisepticemia in Japan. Avian Diseases. 51, 656-662.
- Zhao, X., Yang J., Ju Z., Chang W., Sun S., 2018. Molecular characterization of antimicrobial resistance in Escherichia coli from rabbit farms in Tai'an, China. BioMed Research International. 2018,

Target	Primers sequences	Amplified	Prim. Den	Amplification		Final Extensi	Referenc	
gene		(hn)	Den	Sec Ann Ext		on	03	
		(0p)		den.	Ann.	LAI.	on	
iss	ATGTTATTTTCTGCC	266	94°C	94°C	54°C	72°C	72°C	(Yaguch
	GCTCTG		5min	30sec	30sec	30sec	7	i et al.,
	CTATTGTGAGCAAT						min.	2007)
	ATACCC							
eaeA	ATG CTT AGT GCT	248	94°C	94°C	51°C	72°C	72°C	(Bisi-
	GGT TTA GG		5min	30sec	30sec	30sec	7min	Johnson
	GCC TTC ATC ATT						•	et al.,
	TCG CTT TC							2011)

Table (1): Target genes, oligonucleotides sequences, amplified segment size, and PCR cycling conditions.

Tested antibiotic	Inhibition zone (mm) of isolated <i>E. coli</i>		Interpretive criteria of zone diameter (mm) according (CLSI, 2017)			
	Farm (1)	Farm (2)	Sensitive	intermediate	Resistance	
Ampicillin	15	16	≥17	14–16	≤13	
Tetracycline	0	0	≥15	12–14	≤11	
Norfloxacin	26	28	≥17	13–16	≤12	
Streptomycin	11	14	≥15	12–14	≤11	
Fosfomycin	26	28	≥16	13–15	≤12	
Colistin	13	14	≥11	-	≤10	
Penicillin	15	15	≥29	-	≤28	
erythromycin	0	8-9	≥23	14–22	≤13	
Lincospectin	0	0	≥20	-	≤16	

Table (2): Characterization of antimicrobial resistance pattern in isolated E. coli

Groups	days post <i>E coli</i> infection					
	7 days	15 days	21 days			
G1	921.11±116.05 ^a	987.78±96.92 ^b	1124.44±121.46 ^b			
G2	921.11±116.05 ^a	1147.50±205.27 ^a	1400±180.28 ^a			
G3	936.11±145.53 ^a	1116.67±103.28 ^a	1350 ± 50^{a}			
G4	741.5±208.73 ^b	856.67±120.94 ^b	1210±52.92 ^b			
G5	702.22±121.35 ^b	902.22±121.35 ^b	1150 ± 50^{b}			
G6	912.22±70.89 ^a	980.22 ± 70.89^{b}	1100±50 ^b			

Table (3). The effect of probiotics and herbal oils (fennel and moringa) on body weight of healthy and *E. coli* experimentally infected rabbits at 7^{th} , 15^{th} and 21^{st} days PI. Mean \pm SD.

Means within the same column and carry different small letters see significantly p < 0.05 different G1: non infected non treated (negative control). G2: non infected and treated with probiotic and herbal oils. G3: Infected and treated with probiotic and herbal oils. G4: Infected and treated with probiotic only. G5: Infected and treated with collistin sulfate. G6: infected non treated (positive control).



Figure 1: Detection of E. coli iss (a) and eaeA (b) virulence gene by PCR agarose gelelectrophoresis patternLane L: 100bp DNA ladderLane P: positive controlLane N: negative control



Figure 2: Effects of probiotics, herbal oil and colistin on the total colony count of *E. Coli* in experimentally infected rabbits

G3: Infected and treated with probiotic and herbal oils. G4: Infected and treated with probiotic only. G5: Infected and treated with colistin sulfate. G6: infected non treated (positive control). *P- value < 0.01

Bars carry different small letters sre significantly p < 0.01 different